


anti-Actinin 4

affinity purified rabbit antibody IG701

Lot: A02

data sheet 05/04 — catalog #: 0042-05



immunoGlobe
Antikörpertechnik GmbH

Background information

α -Actinin 4 is an actin-bundling protein of ~100 kDa that is associated with cell motility, endocytosis, and cancer invasion [1, 2]. The α -Actinin family comprises two non-muscle isoforms (α -Actinin 1 and 4) and two skeletal muscle isoforms (α -Actinin 2 and 3), with α -Actinin 2 being also expressed in cardiac muscle [3].

While α -Actinin 4 is almost ubiquitously expressed, particularly high concentrations are found in glomeruli. On the subcellular level it is associated with actin stress fibers, but in certain cells it also localizes to the nucleus [1].

Mutations in the *α -actinin 4* gene cause an autosomal-dominant form of familial focal segmental glomerulosclerosis (FSGS) [4], which is thought to result from a defect in glomerular podocyte function. A point mutation in the *α -actinin 4* gene was found to generate an antigenic peptide that is recognized by autologous cytolytic T lymphocytes (CTL) on a human lung carcinoma [5].

α -Actinin 4 interacts with a variety of proteins, including the ring finger protein BERP [6], the PDZ-LIM protein CLP-36 [7], the hemidesmosomal and cell-cell contact protein BP180 [8], and the tight junction protein MAGI-1 [9]. Moreover, α -Actinin 4 forms a ternary complex with Ca^{2+} /Calmodulin-dependent protein kinase II and Densin-180, a protein of postsynaptic densities in CNS neurons [10]. Ca^{2+} -dependent association of α -Actinin 4 with E3KARP is required for Ca^{2+} -dependent inhibition of the Na^+/H^+ exchanger 3 (NHE3) [11].

Antibody preparation and storage


200 μl of purified antibody in PBS containing 1 mg/ml BSA, 0.01% (w/v) NaN_3 . Antibody concentration: 250 $\mu\text{g}/\text{ml}$. Vials have been overfilled by 10% to ensure complete recovery of the specified amount. Short term storage at 4°C, stable for one year from date of shipment when stored at -20°C. Avoid repeated freezing and thawing! Do not store in "frost-free" freezers.

Antigen

The antibody was raised as described [1] and is directed against a synthetic peptide from the N-terminal part of Actinin 4 (MGDYMAQEDDW) coupled to KLH via a C-terminal cysteine residue. The antibody was affinity purified on the sepharose bound peptide.

Species and Actinin isoform cross-reactivity

By reaction with a set of solid phase bound synthetic peptides, representing the homologous regions of different α -Actinin isoforms, the antibody was shown to be highly specific for Actinin 4 as compared to Actinins 1-3 (Exp. Conditions: 1:10,000 dilution of IG701, enhanced chemoluminescence detection):



# Sequence	Isoform (Species)
1 QTNDYMQPEEDWD	Actinin 1 (human, mouse, rat, chicken)
2 DEDEYMIQEEEDW	Actinin 2 (human, mouse)
3 GGGEYMQEEDWD	Actinin 3 (human, mouse, rat)
4 SMGDYMAQEDDW	<u>Actinin 4 (human, mouse, rat)</u>
5 GMGDYMRPGGDWD	Actinin 4 variant (rat)
6 MDAWYSDEQMDGD	Control (scrambled Actinin 4 peptide)

Species cross-reactivity in immunoblots of total cell or tissue lysates: human, mouse, rat, chinese hamster, pig

Applications

Immunoblotting (0.1-0.25 $\mu\text{g}/\text{ml}$; 1:1,000 - 1:2,500), immuno-fluorescence (1 $\mu\text{g}/\text{ml}$; 1:250), immunoprecipitation (25 $\mu\text{g}/\text{ml}$; 1:10). The recommended dilutions refer to the analysis of mouse cells and tissues with intermediate levels of Actinin 4 expression and must be viewed as approximate.

Positive control

Human platelet protein (500 μg), supplied at 5 mg/ml in SDS sample buffer. Use 5 μl (25 μg) per lane for Western blotting.

(./2)

Related products

- rabbit antiserum M4 to human VASP, 100 µl (catalog # 0010-10)
- Pre-immune serum to M4, 25 µl (catalog # 0013-02)
- affinity purified rabbit antibody IG731 to human VASP, 25 µg (catalog # 0012-02)
- monoclonal antibody IE273 to human VASP, 50 µg (catalog # 0016-05)
- affinity purified rabbit antibody to human LPP, 50 µg (catalog # 0032-05)
- affinity purified rabbit antibody to profilin, 10 µg (catalog # 022-01)

References

1. Honda, K., T. Yamada, R. Endo, Y. Ino, M. Gotoh, H. Tsuda, Y. Yamada, H. Chiba, and S. Hirohashi, *Actinin-4, a novel actin-bundling protein associated with cell motility and cancer invasion*. J Cell Biol, 1998. **140**: 1383-1393.
2. Araki, N., T. Hatae, T. Yamada, and S. Hirohashi, *Actinin-4 is preferentially involved in circular ruffling and macropinocytosis in mouse macrophages: analysis by fluorescence ratio imaging*. J Cell Sci, 2000. **113**: 3329-3340.
3. Critchley, D.R. and G. Flood, *Alpha-Actinins*, in *Guidebook to the Cytoskeletal and Motor Proteins*, T. Kreis and R. Vale, Editors. 1999, Oxford University Press: Oxford. p. 24-27.
4. Kaplan, J.M., S.H. Kim, K.N. North, H. Rennke, L.A. Correia, H.Q. Tong, B.J. Mathis, J.C. Rodriguez-Perez, P.G. Allen, A.H. Beggs, and M.R. Pollak, *Mutations in ACTN4, encoding alpha-actinin-4, cause familial focal segmental glomerulosclerosis*. Nat Genet, 2000. **24**: 251-256.
5. Echchakir, H., F. Mami-Chouaib, I. Vergnon, J.F.

Baurain, V. Karanikas, S. Chouaib, and P.G. Coulie, *A point mutation in the alpha-actinin-4 gene generates an antigenic peptide recognized by autologous cytolytic T lymphocytes on a human lung carcinoma*. Cancer Res, 2001. **61**: 4078-4083.

6. El-Husseini, A.E., D. Kwasnicka, T. Yamada, S. Hirohashi, and S.R. Vincent, *BERP, a novel ring finger protein, binds to alpha-actinin-4*. Biochem Biophys Res Commun, 2000. **267**: 906-911.
7. Vallenius, T., K. Luukko, and T.P. Makela, *CLP-36 PDZ-LIM protein associates with nonmuscle alpha-actinin-1 and alpha-actinin-4*. J Biol Chem, 2000. **275**: 11100-11105.
8. Gonzalez, A.M., C. Otey, M. Edlund, and J.C. Jones, *Interactions of a hemidesmosome component and actinin family members*. J Cell Sci, 2001. **114**: 4197-4206.
9. Patrie, K.M., A.J. Drescher, A. Welihinda, P. Mundel, and B. Margolis, *Interaction of two actin-binding proteins, synaptopodin and alpha-actinin-4, with the tight junction protein MAGI-1*. J Biol Chem, 2002. **277**: 30183-90.
10. Walikonis, R.S., A. Oguni, E.M. Khorosheva, C.J. Jeng, F.J. Asuncion, and M.B. Kennedy, *Densin-180 forms a ternary complex with the (alpha)-subunit of Ca²⁺/calmodulin-dependent protein kinase II and (alpha)-actinin*. J Neurosci, 2001. **21**: 423-433.
11. Kim, J.H., W. Lee-Kwon, J.B. Park, S.H. Ryu, C.H. Yun, and M. Donowitz, *Ca²⁺-dependent inhibition of Na⁺/H⁺ exchanger 3 (NHE3) requires an NHE3-E3KARP-alpha-actinin-4 complex for oligomerization and endocytosis*. J Biol Chem, 2002. **277**: 23714-23724.

for research use only — not for human, in vivo,
diagnostic, therapeutic or other uses